

Antibiotic prophylaxis for Open Fractures



An open (compound) fracture is a fracture with an open wound in the area (even if bone is not visibly protruding). These are often associated with high-energy trauma, and are at increased risk of infection due to surrounding tissue injury and introduction of skin and environmental organisms into the fracture site.

The management of open fractures is focused on preventing infection (through appropriate cleaning of contaminated wound/tissue/bone), antimicrobial prophylaxis, and bone stabilization.

Because the infection risk is dependent on the degree of damage to bone and soft tissues, the choice and duration of antibiotic prophylaxis are based on the classification of open fractures.

CLASSIFICATION OF OPEN FRACTURES

Modified Gustilo-Anderson Grading system

Grade	Description
I	wound < 1 cm, simple fracture pattern, no skin crushing
II	wound 1-10 cm; simple fracture pattern; moderate tissue injury
III A	Wound > 10cm, extensive soft tissue damage but adequate soft-tissue coverage of bone
III B	Wound > 10 cm, extensive soft tissue damage with bone exposure (inadequate soft tissue coverage) or major wound contamination
III C	Wound associated with extensive soft tissue damage with arterial injury requiring repair; concerns for limb preservation

From Samai and Villela (2018) which was adapted from Gustilo and Anderson (1976) and Gustilo, Mendoza and Williams (1984)

ORGANISMS

Gustilo Type I and II: Gram-positive cocci (skin flora):

Gustilo Type III: Gram-positive cocci (skin flora); gram-negatives +/- soil organisms including anaerobes (e.g. *Clostridium sp*)

Timing of treatment initiation: as soon as possible, ideally within 1h of presentation. Delay in administration of antibiotics is associated with poorer outcomes.

Duration of treatment: There is no clear benefit in prolonging prophylactic antibiotics for more than 72h after injury, or 24h after wound closure (whichever comes first)

Open fracture type	Recommended antibiotic prophylaxis *to start as soon as possible, ideally < 1 hour after presentation
Gustilo type I and II	<p>Cefazolin¹ 2 g IV q8h x 24h</p> <p><i>If known MRSA carrier or previous MRSA infection:</i> Cefazolin¹ 2 g IV q8h x 24h AND Vancomycin² 15 mg/kg IV (max 2 g) q12h x 24h</p> <p><i>If severe hypersensitivity reaction to cephalosporins:</i> Vancomycin² 15 mg/kg IV (max 2 g) q12h x 24h</p>
Gustilo type III	<p>Ceftriaxone 2 g IV q24h for up to 24h after wound closure or 72h (<i>whichever comes first</i>)</p> <p><i>If known MRSA carrier or previous MRSA infection:</i> Ceftriaxone 2 g IV q24h AND Vancomycin² 15 mg/kg IV (max 2 g) q12h x 24h after wound closure or 72h (<i>whichever comes first</i>)</p> <p><i>If severe hypersensitivity reaction to cephalosporins:</i> Vancomycin² 15 mg/kg IV (max 2 g) q12h AND Tobramycin³ 5 mg/kg IV (max 400 mg) q24h for up to 24h after wound closure or 72h (<i>whichever comes first</i>)</p>
Soil contamination Fecal contamination Impaired vascularization	<p>Ceftriaxone 2 g IV q24h AND Penicillin⁴ G 4 million units IV q4h for up to 24h after wound closure or 72h (<i>whichever comes first</i>)</p> <p><i>If known MRSA carrier or previous MRSA infection:</i> Add Vancomycin² 15 mg/kg IV (max 2 g) q12h x 24h after wound closure or 72h (<i>whichever comes first</i>)</p> <p><i>If severe hypersensitivity reaction to all beta-lactam antibiotics:</i> Vancomycin² 15 mg/kg IV (max 2 g) q12h AND Tobramycin³ 5 mg/kg IV (max 400 mg) q24h for up to 24h after wound closure or 72h (<i>whichever comes first</i>)</p>
Water contamination (e.g. water sports)	<p>Piperacillin-tazobactam⁵ 4.5 g IV q6h for up to 24h after wound closure or 72h (<i>whichever comes first</i>)</p> <p><i>If known MRSA carrier or previous MRSA infection:</i> Add Vancomycin² 15 mg/kg IV (max 2 g) q12h x 24h after wound closure or 72h (<i>whichever comes first</i>)</p> <p><i>If severe hypersensitivity reaction to penicillin antibiotics:</i> Vancomycin² 15 mg/kg IV (max 2 g) q12h AND Tobramycin³ 5 mg/kg IV (max 400 mg) q24h for up to 24h after wound closure or 72h (<i>whichever comes first</i>)</p>

¹Cefazolin: If weight > 120 kg, increase dose to 3 g IV; cefazolin can be safely given if allergy only to penicillins; if CrCl 10-30 mL/min, give q12h instead of q8h; if CrCl <10 mL/min: 1 g IV q24h

²Vancomycin: If CrCl < 30 mL/min, give q24h instead of q12h; therapeutic drug monitoring not needed if only x 24h

³Tobramycin: If CrCl < 60 mL/min, decrease dose to 2 mg/kg IV; therapeutic drug monitoring not needed if only x 24h

⁴Penicillin: If CrCl 10-50 mL/min: decrease dose to 3 million units; If CrCl < 10 mL/min decrease dose to 2 million units

⁵Piperacillin-tazobactam: If CrCl 20-40 mL/min: 4.5 g IV q8h; CrCl < 20 mL/min: 2.25 g IV q6h; hemodialysis: 2.25 g IV q8h

Drafted by: A. Vintze-Geoffrion (pharmacy resident); reviewed by F. Bourdeau (pharmacy) and M. Semret (ID)

Revised by: Dr Berry (Orthopedic)

Approved by ASP committee on 2021-03-17

Approved by P&T committee on 2022-01-26

References

1. American College of Surgeons - Trauma quality improvement program. ACS TQIP: best practices in management of orthopaedic trauma [Online] 2016. Available : https://www.facs.org/-/media/files/quality-programs/trauma/tqip/ortho_guidelines.ashx

2. Hoff, W. S., Bonadies, J. A., Cachecho, R., & Dorlac, W. C. (2011). East Practice Management Guidelines Work Group: update to practice management guidelines for prophylactic antibiotic use in open fractures. *Journal of Trauma and Acute Care Surgery*, 70(3), 751-754.
3. Garner, M. R., Sethuraman, S. A., Schade, M. A., & Boateng, H. (2020). Antibiotic prophylaxis in open fractures: evidence, evolving issues, and recommendations. *JAAOS-Journal of the American Academy of Orthopaedic Surgeons*, 28(8), 309-315.
4. Rodriguez, L., Jung, H. S., Goulet, J. A., Cicalo, A., Machado-Aranda, D. A., & Napolitano, L. M. (2014). Evidence-based protocol for prophylactic antibiotics in open fractures: improved antibiotic stewardship with no increase in infection rates. *Journal of Trauma and Acute Care Surgery*, 77(3), 400-408.
5. Sudduth, J. D., Moss, J. A., Spittler, C. A., Pham, V. L. H., Jones, L. C., Brown, J. T., & Bergin, P. F. (2020). Open Fractures: Are We Still Treating the Same Types of Infections?. *Surgical Infections*.
6. Hospenthal, D. R., Murray, C. K., Andersen, R. C., Bell, R. B., Calhoun, J. H., Cancio, L. C., ... & Wortmann, G. W. (2011). Guidelines for the prevention of infections associated with combat-related injuries: 2011 update: endorsed by the Infectious Diseases Society of America and the Surgical Infection Society. *Journal of Trauma and Acute Care Surgery*, 71(2), S210-S234.
7. Bratzler, D. W., Dellinger, E. P., Olsen, K. M., Perl, T. M., Auwaerter, P. G., Bolon, M. K., ... & Weinstein, R. A. (2013). Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Surgical infections*, 14(1), 73-156.
8. Chang, Y., Bhandari, M., Zhu, K. L., Mirza, R. D., Ren, M., Kennedy, S. A., ... & Guyatt, G. H. (2019). Antibiotic prophylaxis in the management of open fractures: a systematic survey of current practice and recommendations. *JBJS reviews*, 7(2), e1.
9. Messner, J., Papakostidis, C., Giannoudis, P. V., & Kanakaris, N. K. (2017). Duration of administration of antibiotic agents for open fractures: meta-analysis of the existing evidence. *Surgical Infections*, 18(8), 854-867.
10. Samai, K., & Vilella, A. (2018). Update in Therapeutics: Prophylactic Antibiotics in Open Fractures. *Journal of Trauma Nursing*, 25(2), 83-86.
11. Stennett, C. A., O'Hara, N. N., Sprague, S., Petrisor, B., Jeray, K. J., Leekha, S., ... & FLOW Investigators. (2020). Effect of Extended Prophylactic Antibiotic Duration in the Treatment of Open Fracture Wounds Differs by Level of Contamination. *Journal of Orthopaedic Trauma*, 34(3), 113-120.
12. Bremner, D., Bookstaver, B., Cairns, M., Lindley, K., Durkin, M., Koon, D., & Quidley, A. M. (2017). Impact of body mass index and bacterial resistance in osteomyelitis after antibiotic prophylaxis of open lower-extremity fractures. *Surgical infections*, 18(3), 368-373.
13. Bankhead-Kendall, B., Gutierrez, T., Murry, J., Holland, D., Agrawal, V., Almahmoud, K., ... & Truitt, M. S. (2019). Antibiotics and open fractures of the lower extremity: less is more. *European Journal of Trauma and Emergency Surgery*, 45(1), 125-129.
14. Welborn A.N. When Sticks & Stones Expose Your Bones Will Gentamicin Save You? Evaluating the Need for Expanded Gram-Negative Coverage in Prophylaxis of Grade III Open Fractures. [Online] University of Austin college of Pharmacy. 2018. Available: http://sites.utexas.edu/pharmacotherapy-rounds/files/2018/10/FINAL_Welborn_EGN-OFP_RR-Handout-without-pic-Oct-2018.pdf